

## 16% COOP EGG RATION PELLETS

ITEM NUMBER 270200

### GUARANTEED ANALYSIS

CRUDE PROTEIN, not less than	16.00%
CRUDE FAT, not less than	2.50%
CRUDE FIBER, not more than	9.00%
CALCIUM, not more than	3.80%
CALCIUM, not less than	2.40%
PHOSPHORUS, not less than	.90%

### Ingredients:

Grain Products, Grain Byproducts, Plant Protein Products, Forage Products, Animal Protein Products, Calcium Carbonate, Dicalcium Phosphate, Salt, Vitamin A Supplement, Vitamin D Supplement, Vitamin E Supplement, Riboflavin Supplement, Choline Chloride, Vitamin B-12 Supplement, Niacin, Calcium Pantothenate, Riboflavin Supplement, Manganese Sulphate, Copper Sulphate, Manganese Oxide, Potassium Iodide, Zinc Oxide, Iron Sulfate, and DL Methionine

### Manufactured By

**ALABAMA FARMERS COOPERATIVE, INC.**

**Decatur, Alabama 35601**

### FEEDING DIRECTIONS

Egg Ration 16% protein is a pelleted feed designed to be fed to mature laying hens. The 16% protein level is ideal for mature laying hens and those birds that have free roam. This feed contains high levels of calcium for proper egg shell development. After using this feed for several weeks, if weak egg shells

continue to be a problem, provide another calcium source (oyster Shells) free choice. This feed also contains corn and alfalfa meal. These two products are added to provide xanthophyll. The xanthophylls are a group of naturally occurring pigments which are responsible for the yellow coloration in the egg

yolk, the shanks and feet of the birds, and the yellow color in the fat and skin of birds. This feed also contains reduced levels of salt in an effort to reduce the incidence of wet droppings. These laying feeds have been formulated to include all trace minerals, and vitamins (including B vitamins) which are necessary for egg production. The highest quality ingredients have been used to increase nutrient availability and palatability of this feed to the flock. Supplemental fat has also been added to increase energy to the hen.

### Important Management Practices

Always provide a source of fresh clean water at all times, allow plenty of space for birds to reduce incidence of disease and poor production. Contact an animal nutritionist if you experience prolonged conditions such as weak eggshells, loss of feathers, inability to walk, or blindness. These could all be signs of an improper feeding program.