



PASTURE AID[®] with ALTOSID[®] and CTC (MEDICATED)

For Further Manufacture of Feed

Growing Cattle (over 400 lb.): Increased rate of weight gain, improved feed efficiency and reduction of liver condemnations due to liver abscesses. Calves (250 to 400 lb.): Increased rate of weight gain and improved feed efficiency.
Prevents adult horn fly emergence from manure of treated cattle.

ACTIVE INGREDIENTS*

Methoprene Isopropyl (E,E)-11-methoxy-3,7,11-trimethyl-2,4-dodecadienoate0.01%
Chlortetracycline Hydrochloride448 g/ton

*Each pound contains 45.4 mg of METHOPRENE and 224 mg of CTC

KEEP OUT OF REACH OF CHILDREN CAUTION

SEE REVERSE SIDE FOR ADDITIONAL PRECAUTIONARY
STATEMENTS AND FOR FEEDING DIRECTIONS.

GUARANTEED ANALYSIS

Crude Protein (Min)	12.00%
<small>(This includes not more than 3.5% equivalent protein from non-protein nitrogen).</small>	
Crude Fat (Min)	0.40%
Crude Fiber (Max)	4.50%
Calcium (Min)	10.00%
Calcium (Max)	12.00%
Phosphorus (Min)	1.00%
Salt (Min)	13.00%
Salt (Max)	15.00%
Magnesium (Min)	2.40%
Potassium (Min)	0.50%
Copper (Min)	400 ppm
Iodine (Min)	40 ppm
Magnesium (Min)	1,000 ppm
Selenium	21 ppm
Zinc (Min)	1,600 ppm
Vitamin A (Min)	50,000 IU/lb.
Vitamin D-3 (Min)	10,000 IU/lb.
Vitamin E (Min)	10 IU/lb.

INGREDIENT STATEMENT

Monocalcium Phosphate, Dicalcium Phosphate, Calcium Carbonate, Salt, Cottonseed Meal, Magnesium Limestone, Magnesium Mica, Processed Grain By-products, Urea, Zinc Sulfate, Zinc Oxide, Manganese Sulfate, Manganous Oxide, Copper Sulfate, Ferrous Sulfate, Calcium Iodate, Cobalt Sulfate, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Sodium Selenite, Yeast Culture, Synthetic Red Iron Oxide and Cane Molasses.



NET WEIGHT 50 LB. (22.7 KG)

40094

09/04



Manufactured By Sweetlix, PO Box 8500, Mankato, MN 56002

Contact us at: 1-87SWEETLIX • www.sweetlix.com

**KEEP OUT OF REACH OF CHILDREN
CAUTION
PRECAUTIONARY STATEMENTS**

HAZARDS TO HUMANS

Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water. If irritation persists, get medical attention. Wear long-sleeved shirts and pants; chemical resistant gloves; shoes and sock for protection when handling.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not contaminate water when disposing of equipment wash waters.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.
STORAGE: Store in a dry place in original container. Do not contaminate with pesticides or fertilizers.
CONTAINER DISPOSAL: Completely empty bag into mixing equipment then dispose of empty bag in sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.
PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on-site at an approved waste disposal facility.

MIXING AND FEEDING DIRECTIONS

It is a violation of Federal law to use this product in a manner inconsistent with the labeling.
VMS PASTURE AID[®] WITH ALTOSID[®] AND CTC is formulated to be fed continuously to cattle to provide minerals, vitamins and chlortetracycline daily. Thoroughly mix **VMS PASTURE AID WITH ALTOSID AND CTC** with non-medicated feed at the selected feeding rate. For calves (250-400 lbs.), feed at a rate of 1.8 to 5 oz. per head daily to provide the recommended rate of 25 to 70 mg CTC per day. For growing cattle (over 400 lb.), feed at the rate of 5 oz. per head daily to supply the recommended rate of 70 mg CTC per head per day. Feed at the rate of 5 ounces per head daily to deliver 70 mg CTC and 14 mg Altosid.

Feed Intake (lb/hd/d)	Pasture Aid w/Altosid & CTC (lb/ton)	Non-Medicated Feed (lb/ton)	CTC (mg/hd/day)	Altosid (mg/hd/day)
1.0	625	1375	70	14
5.0	125	1875	70	14
10.0	62.5	1937.5	70	14
20.0	31.25	1968.75	70	14

FLY CONTROL PROGRAM

VMS PASTURE AID[®] WITH ALTOSID[®] AND CTC prevents the development of horn flies in the manure of treated cattle, but is not effective against existing adult flies. When starting a feeding program during the fly season, it may be necessary to use other control measures to reduce the population of existing adult flies. Supplemental fly control measures may be needed in and around buildings to control adult horn flies which breed not only in animal manure but in other decaying matter on the premises. In order to achieve optimum fly control, **VMS PASTURE AID WITH ALTOSID AND CTC** should be used in conjunction with other good management and sanitation practices. Start feeding **VMS PASTURE AID WITH ALTOSID AND CTC** early in the Spring before flies begin to appear and continue feeding throughout the Summer and into the Fall. **ALTOSID** should be fed at a level to provide 22.7 - 45.4 milligrams of **ALTOSID** per 100 pounds of body-weight/month.

Provide clean, fresh water to all animals at all times.

CAUTION: Follow label directions. Consumption of selenium should not exceed 3 mg per head daily.

Altosid[®] is a registered trademark of Wellmark Int'l, Schaumburg, IL

40094

09/04

**KEEP OUT OF REACH OF CHILDREN
CAUTION
PRECAUTIONARY STATEMENTS**

HAZARDS TO HUMANS

Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water. If irritation persists, get medical attention. Wear long-sleeved shirts and pants; chemical resistant gloves; shoes and sock for protection when handling.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not contaminate water when disposing of equipment wash waters.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.
STORAGE: Store in a dry place in original container. Do not contaminate with pesticides or fertilizers.
CONTAINER DISPOSAL: Completely empty bag into mixing equipment then dispose of empty bag in sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.
PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on-site at an approved waste disposal facility.

MIXING AND FEEDING DIRECTIONS

It is a violation of Federal law to use this product in a manner inconsistent with the labeling.
VMS PASTURE AID[®] WITH ALTOSID[®] AND CTC is formulated to be fed continuously to cattle to provide minerals, vitamins and chlortetracycline daily. Thoroughly mix **VMS PASTURE AID WITH ALTOSID AND CTC** with non-medicated feed at the selected feeding rate. For calves (250-400 lbs.), feed at a rate of 1.8 to 5 oz. per head daily to provide the recommended rate of 25 to 70 mg CTC per day. For growing cattle (over 400 lb.), feed at the rate of 5 oz. per head daily to supply the recommended rate of 70 mg CTC per head per day. Feed at the rate of 5 ounces per head daily to deliver 70 mg CTC and 14 mg Altosid.

Feed Intake (lb/hd/d)	Pasture Aid w/Altosid & CTC (lb/ton)	Non-Medicated Feed (lb/ton)	CTC (mg/hd/day)	Altosid (mg/hd/day)
1.0	625	1375	70	14
5.0	125	1875	70	14
10.0	62.5	1937.5	70	14
20.0	31.25	1968.75	70	14

FLY CONTROL PROGRAM

VMS PASTURE AID[®] WITH ALTOSID[®] AND CTC prevents the development of horn flies in the manure of treated cattle, but is not effective against existing adult flies. When starting a feeding program during the fly season, it may be necessary to use other control measures to reduce the population of existing adult flies. Supplemental fly control measures may be needed in and around buildings to control adult horn flies which breed not only in animal manure but in other decaying matter on the premises. In order to achieve optimum fly control, **VMS PASTURE AID WITH ALTOSID AND CTC** should be used in conjunction with other good management and sanitation practices. Start feeding **VMS PASTURE AID WITH ALTOSID AND CTC** early in the Spring before flies begin to appear and continue feeding throughout the Summer and into the Fall. **ALTOSID** should be fed at a level to provide 22.7 - 45.4 milligrams of **ALTOSID** per 100 pounds of body-weight/month.

Provide clean, fresh water to all animals at all times.

CAUTION: Follow label directions. Consumption of selenium should not exceed 3 mg per head daily.

Altosid[®] is a registered trademark of Wellmark Int'l, Schaumburg, IL

40094

09/04