



PEST-A-SIDE MINERAL with RABON® & CTC (MEDICATED)

For Further Manufacture of Feed

For control of fecal flies in the manure of treated cattle. Prevents development of face flies, horn flies, houseflies and stable flies in the manure of treated cattle.

Calves (250 to 400 lb.) Increased rate of weight gain and improved feed efficiency.
 Growing Cattle (over 400 lb.) Increased rate of weight gain, improved feed efficiency and reduction of liver condemnations due to liver abscesses.

ACTIVE DRUG INGREDIENTS

- *Tetrachlorvinphos: (Z) 2-chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate 0.625%
 - *Chlortetracycline Hydrochloride 800 g/ton
- *Each pound contains 2.8 grams of RABON and 400 mg CTC.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

SEE REVERSE SIDE FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND FOR FEEDING DIRECTIONS.

GUARANTEED ANALYSIS

Calcium, (Min)	14.00%
Calcium, (Max)	16.50%
Phosphorus, (Min)	4.00%
Salt, (Min)	19.00%
Salt, (Max)	22.00%
Magnesium, (Min)	5.50%
Potassium, (Min)	0.25%
Cobalt, (Min)	58 ppm
Copper, (Min)	300 ppm
Iodine, (Min)	32 ppm
Iron, (Min)	5500 ppm
Manganese, (Min)	900 ppm
Selenium	26 ppm
Zinc, (Min)	1200 ppm
Vitamin A, (Min)	100,000 IU/lb.
Vitamin D-3, (Min)	25,000 IU/lb.
Vitamin E, (Min)	25 IU/lb.

INGREDIENT STATEMENT

Dicalcium Phosphate, Monocalcium Phosphate, Salt, Calcium Carbonate, Magnesium Limestone, Magnesium Mica, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Cobalt Sulfate, Copper Sulfate, Calcium Iodate, Ferrous Sulfate, Manganese Sulfate, Manganous Oxide, Zinc Sulfate, Zinc Oxide, Sodium Selenite, Synthetic Red Iron Oxide, Yeast Culture and Cane Molasses.



NET WEIGHT 50 LB. (22.7 KG)

40098

09/04



PRECAUTIONARY STATEMENTS

KEEP OUT OF REACH OF CHILDREN

CAUTION:

First Aid: IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger, or if available by administering syrup of ipecac. If person is unconscious, do not give anything by mouth and do not induce vomiting.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

IF ON EYES: Flush with plenty of water. Call a physician if irritation persists.

Note To Physicians And Veterinarians: Symptomatology: Signs of organophosphate toxicity are those of cholinergic overstimulation; hypersalivation, excessive urination, diarrhea and dyspnea (respiratory distress, collapse and death from respiratory muscle paralysis). Treatment: Withdraw product immediately. Clinical includes three categories; muscarinic blocking agents (atropine sulfate), (2) cholinesterase reactivators (atropine and 2-PAM, pralidoxine chloride) and (3) emetics (i.e. syrup of ipecac), cathartics (i.e. vegetable oil) and adsorbents (i.e. charcoal slurry). In case of emergency contact Sweetlix (800) 325-1486.

Hazards to Humans and Domestic Animals: Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction to some individuals. Personal Protective Equipment (PPE): long-sleeved shirt and long pants, shoes and socks, chemical resistant gloves. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Follow manufacturers instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should remove Personal Protective Equipment immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards: This compound is toxic to fish.

Follow label directions. Consumption of selenium should not exceed 3 mg per head per day.

FEEDING AND MANAGEMENT

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

VMS PEST-A-SIDE MINERAL WITH RABON & CTC must be thoroughly mixed with non-medicated feed. For calves (250 - 400 lb.), feed at the rate of 1 to 1.6 oz. per head per day to supply 25 to 40 mg chlortetracycline and 175 to 280 mg Rabon per head per day. For Growing Cattle (over 400 lb.), feed at the rate of 2.8 oz. per head per day to supply the recommended rate of 70 mg chlortetracycline and 490 mg Rabon per head per day.

Feed Intake (lb./hd/d)	PEST-A-SIDE w/RABON & CTC (lb./ton)	Non-Med Feed (lb./ton)	CTC (mg/hd/d)	Rabon (mg/hd/d)
1.0	350	1650	70	490
5.0	70	1930	70	490

WARNING: A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

FLY CONTROL PROGRAM

VMS PEST-A-SIDE MINERAL WITH RABON & CTC prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle, but is not effective against existing adult flies. When starting a feeding program during the fly season, it may be necessary to use other control measures to reduce the population of existing adult flies. In order to achieve fly control, **VMS PEST-A-SIDE MINERAL WITH RABON & CTC** should be used in conjunction with other good management and sanitation practices. Start feeding **VMS PASTURE AID NO.10 WITH ROL AND CTC** early in the spring before flies begin to appear and continue feeding throughout the summer and into fall. **VMS PEST-A-SIDE MINERAL WITH RABON & CTC** may be fed to cattle up to slaughter. The recommended rate for Rabon is 70 mg per 100 lb. bodyweight per day (2.1 grams per 100 lbs. bodyweight per month). The recommended rate for CTC is 25 to 70 mg per head daily for calves 250 to 400 lbs. and 70 mg for calves over 400 lbs.

Provide fresh, clean water to all animals at all times.

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

Storage: Store in a dry place in original container.

Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and Local Authorities, by burning. If burned, stay out of smoke. Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

For additional information, please contact your SWEETLIX representative.

Rabon® is a registered trademark of KMG Chemicals, Houston, TX

40098
09/04

PRECAUTIONARY STATEMENTS

KEEP OUT OF REACH OF CHILDREN

CAUTION:

First Aid: IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger, or if available by administering syrup of ipecac. If person is unconscious, do not give anything by mouth and do not induce vomiting.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

IF ON EYES: Flush with plenty of water. Call a physician if irritation persists.

Note To Physicians And Veterinarians: Symptomatology: Signs of organophosphate toxicity are those of cholinergic overstimulation; hypersalivation, excessive urination, diarrhea and dyspnea (respiratory distress, collapse and death from respiratory muscle paralysis). Treatment: Withdraw product immediately. Clinical includes three categories; muscarinic blocking agents (atropine sulfate), (2) cholinesterase reactivators (atropine and 2-PAM, pralidoxine chloride) and (3) emetics (i.e. syrup of ipecac), cathartics (i.e. vegetable oil) and adsorbents (i.e. charcoal slurry). In case of emergency contact Sweetlix (800) 325-1486.

Hazards to Humans and Domestic Animals: Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction to some individuals. Personal Protective Equipment (PPE): long-sleeved shirt and long pants, shoes and socks, chemical resistant gloves. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Follow manufacturers instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should remove Personal Protective Equipment immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards: This compound is toxic to fish.

Follow label directions. Consumption of selenium should not exceed 3 mg per head per day.

FEEDING AND MANAGEMENT

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

VMS PEST-A-SIDE MINERAL WITH RABON & CTC must be thoroughly mixed with non-medicated feed. For calves (250 - 400 lb.), feed at the rate of 1 to 1.6 oz. per head per day to supply 25 to 40 mg chlortetracycline and 175 to 280 mg Rabon per head per day. For Growing Cattle (over 400 lb.), feed at the rate of 2.8 oz. per head per day to supply the recommended rate of 70 mg chlortetracycline and 490 mg Rabon per head per day.

Feed Intake (lb./hd/d)	PEST-A-SIDE w/RABON & CTC (lb./ton)	Non-Med Feed (lb./ton)	CTC (mg/hd/d)	Rabon (mg/hd/d)
1.0	350	1650	70	490
5.0	70	1930	70	490

WARNING: A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

FLY CONTROL PROGRAM

VMS PEST-A-SIDE MINERAL WITH RABON & CTC prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle, but is not effective against existing adult flies. When starting a feeding program during the fly season, it may be necessary to use other control measures to reduce the population of existing adult flies. In order to achieve fly control, **VMS PEST-A-SIDE MINERAL WITH RABON & CTC** should be used in conjunction with other good management and sanitation practices. Start feeding **VMS PASTURE AID NO.10 WITH ROL AND CTC** early in the spring before flies begin to appear and continue feeding throughout the summer and into fall. **VMS PEST-A-SIDE MINERAL WITH RABON & CTC** may be fed to cattle up to slaughter. The recommended rate for Rabon is 70 mg per 100 lb. bodyweight per day (2.1 grams per 100 lbs. bodyweight per month). The recommended rate for CTC is 25 to 70 mg per head daily for calves 250 to 400 lbs. and 70 mg for calves over 400 lbs.

Provide fresh, clean water to all animals at all times.

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

Storage: Store in a dry place in original container.

Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and Local Authorities, by burning. If burned, stay out of smoke. Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

For additional information, please contact your SWEETLIX representative.

Rabon® is a registered trademark of KMG Chemicals, Houston, TX

40098
09/04