



PASTURE AID[®]

Fly Control Mineral

with RABON[®] Oral Larvacide

For control of fecal flies in the manure of treated cattle.
Prevents development of face flies, horn flies, house flies and stable flies in the manure of treated cattle.

ACTIVE INGREDIENT:	BY WEIGHT:
* Tetrachlorvinphos: (Z)2-chloro-1 (2,4,5-trichlorophenyl) vinyl dimethyl phosphate	0.494%
** OTHER INGREDIENTS	99.506%
TOTAL	100.000%

* Rabon[®] insecticide - contains 2.24grams of RABON per pound.

** Refers only to ingredients which are not larvicidal

EPA REG NO. 73600-3

EPA EST. NO. 73600-AL-01

KEEP OUT OF REACH OF CHILDREN

CAUTION

See reverse for additional precautionary statements
and for feeding directions.

GUARANTEED ANALYSIS

Calcium (Min)	18.40%
Calcium (Max)	22.00%
Phosphorus (Min)	2.00%
Salt (Min)	20.00%
Salt (Max)	24.00%
Magnesium (Min)	4.00%
Potassium (Min)	0.10%
Copper (Min)	200 ppm
Iodine (Min)	48 ppm
Selenium	18 ppm
Zinc (Min)	800 ppm
Vitamin A (Min)	50,000 IU/lb.
Vitamin D-3 (Min)	12,500 IU/lb.
Vitamin E (Min)	12.5 IU/lb.

INGREDIENT STATEMENT

Defluorinated Phosphate, Calcium Carbonate, Magnesium Limestone, Salt, Processed Grain By-products, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Ferrous Sulfate, Manganous Oxide, Managese Sulfate, Zinc Oxide, Zinc Sulfate, Cobalt Sulfate, Copper Sulfate, Calcium Iodate, Sodium Selenite, Plant Protein Products, Synthetic Red Iron Oxide and Molasses Products.



NET WEIGHT 50 LB. (22.7 KG)

Manufactured By Sweetlix, PO Box 8500, Mankato, MN 56002

Contact us at: 1-87SWEETLIX • www.sweetlix.com

09/04

FIRST AID

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 to 2 glasses of water and induce vomiting by touching back of throat with finger, or if available by administering syrup of ipecac. If person is unconscious, do not give by mouth and do not induce vomiting.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

IF IN EYES: Flush with plenty of water. Call a physician if irritation persists.

NOTE TO PHYSICIANS AND VETERINARIANS

Symptomatology: Signs of organophosphate toxicity are those of cholinergic over stimulation; hypersalivation, excessive urination, diarrhea and dyspnea (respiratory distress, collapse and death from respiratory muscle paralysis).

Treatment: Withdraw product immediately. Clinical includes 3 categories; musarinic blocking agents (atropine sulfate), (2) cholinesterase reactivators (atropine and 2-PAM, pralidoxine chloride) and (3) emetics (i.e. syrup of ipecac), cathartics (i.e. vegetable oil) and absorbents (i.e. charcoal slurry). In case of emergency, contact Sweetlix LLC (800) 325-1486.

PRECAUTIONARY STATEMENTS

CAUTION:

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction to some individuals. Personal Protective Equipment (PPE): long-sleeved shirt and long pants, shoes and socks, chemical resistant gloves. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Follow manufacturers instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards: This compound is toxic to fish.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

V.M.S. PASTURE AID® FLY CONTROL MINERAL with Rabon® Oral Larvicide is formulated to be fed continuously on a free choice basis to cattle to provide minerals, vitamins and Rabon® daily. Feed supplies (pasture or forage) and feeder locations should be adequate to assure consumption in accordance with the intake levels below:

CATTLE WEIGHT (Lb.)	RABON® SUPPLEMENT Daily Intake		RABON® Daily Intake (Mg)
	(Oz.)	(Lb.)	
500	2.5	0.15625	350
600	3.0	0.18750	420
700	3.5	0.21875	490
800	4.0	0.25000	560
900	4.5	0.28125	630
1000	5.0	0.31250	700
1100	5.5	0.34375	770
1200	6.0	0.37500	840

If cattle consume more or less than these levels, feeders should be moved further from or closer to the general resting and watering area. If cattle still fail to consume the correct amount, discontinue feeding this free choice supplement. Feed V.M.S. PASTURE AID FLY CONTROL MINERAL with Rabon® Oral Larvicide as the only FREE CHOICE source of salt, other minerals and vitamins.

FLY CONTROL PROGRAM

V.M.S. Pasture Aid® Fly Control Mineral with Rabon Oral Larvicide prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle, but is not effective against existing adult flies. When starting a feeding program during the fly season, it may be necessary to use other control measures to reduce the population of existing adult flies. Supplemental fly control measures may be needed in and around buildings to control adult house flies and stable flies which breed not only in animal manure but in other decaying matter on the premises. In order to achieve optimum fly control, V.M.S. Pasture Aid Fly Control Mineral with Rabon Oral Larvicide should be used in conjunction with other good management and sanitation practices. Start feeding V.M.S. Pasture Aid Fly Control Mineral with Rabon Oral Larvicide early in the spring before flies begin to appear and continue feeding throughout the summer and into fall. V.M.S. Pasture Aid Fly Control Mineral with Rabon Oral Larvicide may be fed to cattle up to slaughter. Rabon Oral Larvicide should be fed at a level to provide 70 mg Rabon per 100 lb. of bodyweight per day (2.1 grams per 100 lbs. bodyweight per month).

Provide fresh, clean water to all animals at all times.

Follow label directions. Consumption of selenium should not exceed 3 mg per head per day.

Rabon® is a registered trademark of KMG Chemicals, Houston, TX

40099

09/04

FIRST AID

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 to 2 glasses of water and induce vomiting by touching back of throat with finger, or if available by administering syrup of ipecac. If person is unconscious, do not give by mouth and do not induce vomiting.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

IF IN EYES: Flush with plenty of water. Call a physician if irritation persists.

NOTE TO PHYSICIANS AND VETERINARIANS

Symptomatology: Signs of organophosphate toxicity are those of cholinergic over stimulation; hypersalivation, excessive urination, diarrhea and dyspnea (respiratory distress, collapse and death from respiratory muscle paralysis).

Treatment: Withdraw product immediately. Clinical includes 3 categories; musarinic blocking agents (atropine sulfate), (2) cholinesterase reactivators (atropine and 2-PAM, pralidoxine chloride) and (3) emetics (i.e. syrup of ipecac), cathartics (i.e. vegetable oil) and absorbents (i.e. charcoal slurry). In case of emergency, contact Sweetlix LLC (800) 325-1486.

PRECAUTIONARY STATEMENTS

CAUTION:

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction to some individuals. Personal Protective Equipment (PPE): long-sleeved shirt and long pants, shoes and socks, chemical resistant gloves. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Follow manufacturers instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards: This compound is toxic to fish.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

V.M.S. PASTURE AID® FLY CONTROL MINERAL with Rabon® Oral Larvicide is formulated to be fed continuously on a free choice basis to cattle to provide minerals, vitamins and Rabon® daily. Feed supplies (pasture or forage) and feeder locations should be adequate to assure consumption in accordance with the intake levels below:

CATTLE WEIGHT (Lb.)	RABON® SUPPLEMENT Daily Intake		RABON® Daily Intake (Mg)
	(Oz.)	(Lb.)	
500	2.5	0.15625	350
600	3.0	0.18750	420
700	3.5	0.21875	490
800	4.0	0.25000	560
900	4.5	0.28125	630
1000	5.0	0.31250	700
1100	5.5	0.34375	770
1200	6.0	0.37500	840

If cattle consume more or less than these levels, feeders should be moved further from or closer to the general resting and watering area. If cattle still fail to consume the correct amount, discontinue feeding this free choice supplement. Feed V.M.S. PASTURE AID FLY CONTROL MINERAL with Rabon® Oral Larvicide as the only FREE CHOICE source of salt, other minerals and vitamins.

FLY CONTROL PROGRAM

V.M.S. Pasture Aid® Fly Control Mineral with Rabon Oral Larvicide prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle, but is not effective against existing adult flies. When starting a feeding program during the fly season, it may be necessary to use other control measures to reduce the population of existing adult flies. Supplemental fly control measures may be needed in and around buildings to control adult house flies and stable flies which breed not only in animal manure but in other decaying matter on the premises. In order to achieve optimum fly control, V.M.S. Pasture Aid Fly Control Mineral with Rabon Oral Larvicide should be used in conjunction with other good management and sanitation practices. Start feeding V.M.S. Pasture Aid Fly Control Mineral with Rabon Oral Larvicide early in the spring before flies begin to appear and continue feeding throughout the summer and into fall. V.M.S. Pasture Aid Fly Control Mineral with Rabon Oral Larvicide may be fed to cattle up to slaughter. Rabon Oral Larvicide should be fed at a level to provide 70 mg Rabon per 100 lb. of bodyweight per day (2.1 grams per 100 lbs. bodyweight per month).

Provide fresh, clean water to all animals at all times.

Follow label directions. Consumption of selenium should not exceed 3 mg per head per day.

Rabon® is a registered trademark of KMG Chemicals, Houston, TX

40099

09/04