



PASTURE AID BAR FLY BLOCK #10



A FREE CHOICE SUPPLEMENT BLOCK FOR BEEF CATTLE

For control of fecal flies in the manure of treated cattle. Prevents the development of face flies, horn flies, stable flies and house flies in the manure of treated cattle.

ACTIVE INGREDIENT:*

- Tetrachlorvinphos: (Z) 2-chloro-1 (2,4,5-trichlorophenyl) vinyl dimethyl phosphate 0.30%
- OTHER INGREDIENTS ** 99.70%
- TOTAL 100.00%

*Each pound contains 1.36g Rabon®
 **Refers only to ingredients which are not larvicidal

GUARANTEED ANALYSIS

- Crude Protein (Min) 10.00%
(This includes not more than 3.5% equivalent protein from non-protein nitrogen).
- Crude Fat (Min) 0.50%
- Crude Fiber (Max) 6.50%
- Calcium (Min) 10.50%
- Calcium (Max) 12.50%
- Phosphorus (Min) 1.00%
- Salt (Min) 10.00%
- Salt (Max) 12.00%
- Magnesium (Min) 4.00%
- Potassium (Min) 0.50%
- Copper (Min) 400 ppm
- Iodine (Min) 40 ppm
- Selenium (Min) 13 ppm
- Zinc (Min) 1,600 ppm
- Vitamin A (Min) 20,000 IU/lb.
- Vitamin D-3 (Min) 5,000 IU/lb.
- Vitamin E (Min) 5 IU/lb.

FEED INGREDIENTS

Magnesium Limestone, Salt, Defluorinated Phosphate, Cane Molasses, Soybean Hulls, Dehydrated Alfalfa, Cottonseed Meal, Lignin Sulfonate, Urea, Potassium Sulfate, Magnesium Sulfate, Processed Grain By-Products, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Ferrous Sulfate, Manganese Sulfate, Manganous Oxide, Zinc Sulfate, Zinc Oxide, Cobalt Sulfate, Calcium Iodate, Copper Sulfate, Yeast Culture, Sodium Selenite and Calcium Carbonate.

Keep Out of Reach of Children

Manufactured By Sweetlix, PO Box 8500, Mankato, MN 56002
 Contact us at: 1-877SWEETLIX • www.sweetlix.com

CAUTION

First Aid	
IF SWALLOWED	Call a Poison Control Center or physician immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF ON SKIN	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
IF IN EYES	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER: In case of emergency, contact Sweetlix at (800) 325-1486. For information on this pesticide product (including health concerns, medical emergencies or pesticide incidents), call the National Pesticide Information Center at (800) 858-7378.	
NOTE TO PHYSICIANS: This product is a cholinesterase inhibitor. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.	

PRECAUTIONARY STATEMENTS
 HAZARDS TO HUMANS AND DOMESTIC ANIMALS:

CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction to some individuals.

PERSONAL PROTECTIVE EQUIPMENT (PPE): Long-sleeved shirt and long pants, shoes and socks, chemical resistant gloves.
 USER SAFETY RECOMMENDATIONS: Users should remove Personal Protective Equipment immediately after handling this product. Wash the outside of gloves before removing.

As soon as possible, wash thoroughly and change into clean clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
 USER SAFETY REQUIREMENTS: Follow manufacturers instructions for cleaning/ maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENVIRONMENTAL HAZARDS: This compound is toxic to fish.

DIRECTIONS FOR USE:

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- (1) Provide one block per 10-15 head of cattle. Cattle should consume an average of 1.67 lb. of the block per 100 pounds of body weight per month. The recommended average intake is 70 mg Rabon per hundred pounds of body weight (2.1 g per 100 lbs. of body weight per month). Do not feed other free choice sources of salt. Animals should not be starved for minerals before use of this block.
 - (2) If consumption is above recommended amounts, reduce the number of feeding locations or relocate. If below, increase the number of feeding locations or relocate.
 - (3) Rabon is not effective against existing adult flies. When starting a feeding program during the fly season, it may be necessary to use other control measures to reduce the population of existing adult flies.
 - (4) Consumption of Selenium should not exceed 3 mg daily.

STORAGE AND DISPOSAL: Do not contaminate water, food, or feed by storage and disposal.

Storage: Store in a dry place in original container.
 Pesticide Disposal: Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.
 Container Disposal: Dispose of empty container in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

For additional information, please contact your Sweetlix representative.

02/05



Rabon® is a registered trademark of KMG Chemicals, Houston, TX

40109

NET WEIGHT 33.3 LB. (15.1 KG) EPA REG NO. 73600-5 • EPA EST. NO. 73600-AL-01