



MGA/BOVATEC[®] PREMIX

(MEDICATED)

For Further Manufacture of Feed

For increased rate of weight gain, improved feed efficiency and suppression of Estrus (heat) in heifers fed in confinement for slaughter.

ACTIVE INGREDIENTS

MELENGESTROL ACETATE0.000176%
BOVATEC (as Lasalocid Sodium)1,440 g/ton
Each lb. contains 0.80 mg melengestrol acetate and 720 mg lasalocid.

GUARANTEED ANALYSIS

Calcium (Min) 27.00%
Calcium (Max) 32.00%
Phosphorus (Min) 6.00%
Salt (Min) None Added
Magnesium (Min) 2.75%
Potassium (Min) 0.50%
Sulfur (Min) 0.90%
Cobalt (Min) 10 ppm
Copper (Min) 400 ppm
Iodine (Min) 40 ppm
Iron (Min) 4,000 ppm
Manganese (Min) 1,200 ppm
Selenium 12 ppm
Zinc (Min) 1,600 ppm
Vitamin A (Min) 100,000 IU/lb.
Vitamin D-3 (Min) 25,000 IU/lb.
Vitamin E (Min) 25 IU/lb.

INGREDIENT STATEMENT

Dicalcium Phosphate, Monocalcium Phosphate, Calcium Carbonate, Magnesium Oxide, Magnesium Sulfate, Potassium Sulfate, Cobalt Sulfate, Copper Sulfate, Calcium Iodate, Ferrous Sulfate, Manganese Sulfate, Manganous Oxide, Sodium Selenite, Zinc Sulfate, Zinc Oxide, Vitamin A Supplement, D-3 Supplement, Vitamin E Supplement, Yeast Culture and Mineral Oil.

Important: Only for use in heifers being fed in confinement for slaughter.



NET WEIGHT 50 LB. (22.7 KG)

40124

09/04



Manufactured By Sweetlix, PO Box 8500, Mankato, MN 56002
Contact us at: 1-87SWEETLIX • www.sweetlix.com

MIXING DIRECTIONS

Thoroughly mix the levels of VMS MGA/Bovatec Premix indicated below with non-medicated grain and/or roughages to provide the drug levels indicated:

Feed Intake (lbs)	MGA/Bovatec Premix (lbs/ton)	Non-Medicated Feed (lbs/ton)	Bovatec (mg/hd/day)	MGA (mg/hd/day)
5.0	125	1875	225	0.25
7.5	125	1875	225	0.25
5.0	200	1800	360	0.40
10.0	100	1900	360	0.40
20.0	50	1950	360	0.40

For the stated drug claims heifers should receive not less than 100 mg nor more than 360 mg Lasalocid per head daily and 0.25 to 0.50 mg MGA per head daily. Eight (8) ounces of this premix will deliver 360 mg Lasalocid, 0.4 mg MGA and 2.7 mg selenium.

Provide clean, fresh water to animals at all times.

CAUTION: USE AS DIRECTED

The safety of Lasalocid in unapproved species has not been established. Do not allow horses or other equines access to Lasalocid as ingestion may be fatal. Feeding undiluted or mixing errors resulting in high concentrations of Lasalocid could be fatal to cattle; it must be mixed thoroughly in feeds before use. Do not feed undiluted.

CAUTION: Follow label directions. Consumption of selenium should not exceed 3 mg per head daily.

For additional information, please contact your SWEETLIX representative.

Bovatec[®] is a registered trademark of Alpharma Animal Health Division, Fort Lee, NJ

MIXING DIRECTIONS

Thoroughly mix the levels of VMS MGA/Bovatec Premix indicated below with non-medicated grain and/or roughages to provide the drug levels indicated:

Feed Intake (lbs)	MGA/Bovatec Premix (lbs/ton)	Non-Medicated Feed (lbs/ton)	Bovatec (mg/hd/day)	MGA (mg/hd/day)
5.0	125	1875	225	0.25
7.5	125	1875	225	0.25
5.0	200	1800	360	0.40
10.0	100	1900	360	0.40
20.0	50	1950	360	0.40

For the stated drug claims heifers should receive not less than 100 mg nor more than 360 mg Lasalocid per head daily and 0.25 to 0.50 mg MGA per head daily. Eight (8) ounces of this premix will deliver 360 mg Lasalocid, 0.4 mg MGA and 2.7 mg selenium.

Provide clean, fresh water to animals at all times.

CAUTION: USE AS DIRECTED

The safety of Lasalocid in unapproved species has not been established. Do not allow horses or other equines access to Lasalocid as ingestion may be fatal. Feeding undiluted or mixing errors resulting in high concentrations of Lasalocid could be fatal to cattle; it must be mixed thoroughly in feeds before use. Do not feed undiluted.

CAUTION: Follow label directions. Consumption of selenium should not exceed 3 mg per head daily.

For additional information, please contact your SWEETLIX representative.

Bovatec[®] is a registered trademark of Alpharma Animal Health Division, Fort Lee, NJ