



LIVESTOCK MINERAL • VITAMIN SUPPLEMENT

# CTC -2000 (MEDICATED)

## For Further Manufacture of Feed

Growing Cattle over 400 lbs: For increased rate of weight gain and improved feed efficiency and reduction of liver condemnations due to liver abscesses.

Beef Cattle: For control of bacterial pneumonia associated with shipping fever complex caused by *Pasteurella* spp. Susceptible to chlortetracycline.

Beef Cattle up to 1000 lbs: For control of active anaplasmosis caused by *Anaplasma marginale* susceptible to chlortetracycline.

### ACTIVE DRUG INGREDIENT:

Chlortetracycline ..... 4 g/ton  
(Each lb. contains 2000 mg CTC)

### GUARANTEED ANALYSIS

Calcium, (Min) .....	16.50%
Calcium, (Max) .....	19.80%
Phosphorus, (Min) .....	5.50%
Salt, (Min) .....	11.00%
Salt, (Max) .....	13.00%
Magnesium, (Min) .....	6.00%
Potassium, (Min) .....	1.50%
Sulfur, (Min) .....	2.00%
Cobalt, (Min) .....	19 ppm
Copper, (Min) .....	1000 ppm
Iodine, (Min) .....	100 ppm
Manganese, (Min) .....	2600 ppm
Selenium .....	26 ppm
Zinc, (Min) .....	4000 ppm
Vitamin A, (Min) .....	160,000 IU/LB.
Vitamin D-3, (Min) .....	40,000 IU/LB.
Vitamin E, (Min) .....	25 IU/LB.

### INGREDIENT STATEMENT

Dicalcium Phosphate, Monocalcium Phosphate, Salt, Magnesium Limestone, Defluorinated Phosphate, Magnesium Sulfate, Potassium Sulfate, Calcium Carbonate, Zinc Oxide, Copper Sulfate, Manganous Oxide, Zinc Sulfate, Ferrous Sulfate, Manganese Sulfate, Calcium Iodate, Yeast Culture, Cobalt Sulfate, Sodium Selenite, Magnesium Oxide, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Synthetic Red Iron Oxide And Cane Molasses.

See reverse side for feeding directions and precautionary statements.



NET WEIGHT 50 LB. (22.7 KG)

# 40143

09/04



Manufactured By Sweetlix, PO Box 8500, Mankato, MN 56002

Contact us at: 1-87SWEETLIX • www.sweetlix.com

## MIXING DIRECTIONS

VMS CTC-2000 should be thoroughly mixed with non-medicated grain or roughages at the rates indicated below to form a medicated feed.

Lbs. of VMS CTC-2000 per ton	Lbs. of Non-Medicated Feed per Ton	Lbs. of Feed to Supply mg/hd/day		
		70	350	500
50	1950	1.4	7	10
100	1900	0.7	3.5	5
150	1850	0.5	2.3	3.3

*For anaplasmosis control:* Cattle 700 to 1000 lbs should receive 0.5 mg CTC per pound of bodyweight per day; thus a 1000 lb. cow would require 500 mg. Cattle under 700 lbs. should receive 350 mg CTC per head per day.

*For bacterial pneumonia control:* Beef cattle should receive 350 mg CTC per head per day.

*For improved weight gain and feed efficiency & reduction of liver condemnations:* Growing cattle over 400 lbs. should receive 70 mg CTC per head per day.

Provide clean, fresh water to all animals at all times.

### WARNING:

**DISCONTINUE USE 48 HOURS BEFORE SLAUGHTER.**

A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

### CAUTION: USE AS DIRECTED.

Not for use in lactating dairy cattle. Consumption of selenium should not exceed 3 mg. daily. Four (4) ounces of this product will supply 3 mg. of Selenium.

For additional information, please contact your SWEETLIX representative.

## MIXING DIRECTIONS

VMS CTC-2000 should be thoroughly mixed with non-medicated grain or roughages at the rates indicated below to form a medicated feed.

Lbs. of VMS CTC-2000 per ton	Lbs. of Non-Medicated Feed per Ton	Lbs. of Feed to Supply mg/hd/day		
		70	350	500
50	1950	1.4	7	10
100	1900	0.7	3.5	5
150	1850	0.5	2.3	3.3

*For anaplasmosis control:* Cattle 700 to 1000 lbs should receive 0.5 mg CTC per pound of bodyweight per day; thus a 1000 lb. cow would require 500 mg. Cattle under 700 lbs. should receive 350 mg CTC per head per day.

*For bacterial pneumonia control:* Beef cattle should receive 350 mg CTC per head per day.

*For improved weight gain and feed efficiency & reduction of liver condemnations:* Growing cattle over 400 lbs. should receive 70 mg CTC per head per day.

Provide clean, fresh water to all animals at all times.

### WARNING:

**DISCONTINUE USE 48 HOURS BEFORE SLAUGHTER.**

A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

### CAUTION: USE AS DIRECTED.

Not for use in lactating dairy cattle. Consumption of selenium should not exceed 3 mg. daily. Four (4) ounces of this product will supply 3 mg. of Selenium.

For additional information, please contact your SWEETLIX representative.