

# Foragemax 25% Supplement Block

FOR MATURE BEEF CATTLE ON PASTURE

Manufactured with  
**En-Pro-AI**<sup>®</sup>  
TECHNOLOGY

## GUARANTEED ANALYSIS

Crude Protein (Min) . . . . .	25.00%
(This includes not more than 19.5% equivalent protein from non-protein nitrogen.)	
Crude Fat (Min) . . . . .	5.00%
Crude Fiber (Max) . . . . .	9.00%
Calcium (Min) . . . . .	0.50%
Calcium (Max) . . . . .	1.00%
Phosphorus (Min) . . . . .	0.06%
Salt (Min) . . . . .	4.75%
Salt (Max) . . . . .	5.25%
Magnesium (Min) . . . . .	3.00%
Potassium (Min) . . . . .	2.00%
Selenium (Min) . . . . .	3.4 ppm
Vitamin A (Min) . . . . .	10,000 IU/lb.
Vitamin D-3 . . . . .	2,500 IU/lb.
Vitamin E (Min) . . . . .	2.5 IU/lb

## INGREDIENTS

Cane Molasses, Peanut Hulls, Hydrolyzed Feathermeal, Urea, Hydrolyzed Vegetable Oil, Defluorinated Phosphate, Magnesium Oxide, Salt, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Ferrous Carbonate, Calcium Iodate, Zinc Oxide, Manganous Oxide, Copper Oxide, Cobalt Carbonate, Sodium Selenite, Calcium Carbonate and Mineral Oil.



Contact us at:

**1-87SWEETLIX**  
**www.sweetlix.com**

Manufactured By Sweetlix, PO Box 8500, Mankato, MN 56002

**NET WEIGHT 200 LB. (90.90 KG)**

## FEEDING AND MANAGEMENT

1. Feed one block per 20 head of cattle. (Timid and smaller animals need their share).
2. Place the blocks where animals congregate (loafing, grazing, feeding and watering areas).
3. Feed blocks continuously along with plentiful sources of good quality forage and clean, fresh water.
4. Cattle will consume 1-2 pounds per head daily based upon a 1,000 pound animal. Two (2) pounds of this product will supply 3 mg of selenium.
5. It is also recommended that a Sweetlix free choice mineral be fed at the same time this product is being used for optimum utilization of available roughages.

**Important:** Consumption of this block will vary depending upon the condition of the animals, quality and quantity of forages being fed, seasonal weather conditions and most importantly, the feeding location of the blocks with respect to loafing, grazing, feeding and watering area.

**Introduction Period:** During the first week of use, place the blocks approximately 50-75 feet from loafing and/or watering areas. As cattle become acclimated to the presence of the blocks, some repositioning may be necessary to achieve proper consumption. Move blocks closer to increase consumption; further away to decrease consumption. **DO NOT PROVIDE STARVED CATTLE FREE ACCESS TO THIS BLOCK.**

**Important:** If over consumption occurs for more than two weeks and repositioning of the blocks does not correct the situation, remove blocks and evaluate overall feeding program.

**CAUTION: USE AS DIRECTED. DO NOT FEED TO HORSES.**

Consumption of selenium should not exceed 3 mg per head per day.

08/04



**41541**