

24% Poured Block

(For Ruminants Only)

For Mature Beef Cattle on Pasture

Forage Supplement

GUARANTEED ANALYSIS:

Crude Protein (Min).....	24.00%
(This includes not more than 18% equivalent from non-protein nitrogen.)	
Crude Fat (Min).....	5.00%
Crude Fiber (Max).....	6.00%
Calcium (Min).....	0.50%
Calcium (Max).....	1.00%
Phosphorus (Min).....	0.08%
Salt (Min).....	2.50%
Salt (Max).....	3.00%
Magnesium (Min).....	3.00%
Potassium (Min).....	2.00%
Selenium.....	3.4 ppm
Vitamin A (Min).....	15,000 IU/lb.
Vitamin D-3 (Min).....	3,750 IU/lb.
Vitamin E (Min).....	3.75 IU/lb.

INGREDIENTS:

Cane Molasses, Peanut Hulls, Urea, Magnesium Oxide, Salt, Hydrolyzed Vegetable Oil, Hydrolyzed Feather Meal, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Ferrous Carbonate, Calcium Iodate, Zinc Oxide, Manganous Oxide, Copper Oxide, Cobalt Carbonate, Calcium Iodate and Sodium Selenite.

Important: Consumption of this block will vary depending upon the condition of the animals, quality and quantity of forages being fed, seasonal weather conditions and most importantly, the feeding location of the blocks with respect to loafing, grazing, feeding and watering areas.

CAUTION: Use As Directed. DO NOT FEED TO HORSES.

Consumption of selenium should not exceed 3 mg per head per day.

Important: If over consumption occurs for more than two weeks and repositioning of the blocks does not correct the situation, remove blocks and evaluate overall feeding program.

FEEDING AND MANAGEMENT

1. Feed one block per 20 head of cattle. (Timid and smaller animals need their share.)
2. Place the blocks where animals congregate (loafing, grazing, feeding and watering areas).
3. Feed blocks continuously along with plentiful sources of good quality forage and clean, fresh water.
4. Cattle will consume 1-2 pounds per head daily based upon a 1,000 pound animal. Two pounds of this product will supply 3 mg of selenium.
5. It is also recommended that a Sweetlix free choice mineral be fed with the SWEETLIX 24% POURED BLOCK for optimum utilization of roughages.

Introduction Period: During the first week of use, place the blocks approximately 50-75 feet from loafing and/or watering areas. As cattle become acclimated to the presence of the blocks, some repositioning may be necessary to achieve proper consumption. Move blocks closer to increase consumption; further away to decrease consumption.

DO NOT PROVIDE STARVED CATTLE FREE ACCESS TO THIS BLOCK.

For additional information, please contact your SWEETLIX representative.

08/04

NET WEIGHT: 50 LBS. (22.7 KG)



Manufactured with
En-Pro-Al[®]
TECHNOLOGY