

CO-OP SHEEP AND GOAT FEED

W/DECCOX/A/N

Item number 293040

MEDICATED

For the prevention of coccidiosis in young sheep caused by *E. christenseni*, and *E. ninakohlyakimovae*.

ACTIVE DRUG INGREDIENT

Decoquinatate 15 GM/TON

Warning:

Do not feed to sheep or goats producing milk for food

Directions:

Feed continuously at a rate to provide Decoquinatate at a daily dose of 22.7mg/100 lbs. of body weight and feed for at least 28 days during exposure or when experience indicates that coccidiosis is likely. This feed contains 7.5mg of Decoquinatate per lb. Of complete feed.

GUARANTEED ANALYSIS:

Crude Protein, not less than	15.00%
Crude Fat, not less than	2.00%
Crude Fiber, not more than	15.00%

INGREDIENTS:

Processed Grain By-Products, Plant Protein Products, Grain Products, Roughage Products, Calcium Carbonate, Salt, Vitamin A Supplement, Vitamin D Supplement, Vitamin E Supplement, Magnous Oxide, Magnous Sulfate, Zinc Sulfate, Zinc Oxide, Ferrous Carbonate, Calcium Iodate, Cobalt Carbonate, Sodium Selenium , Manganese Sulphate, Copper Sulfate, Potassium Iodide, and Iron Sulphate

Manufactured By

ALABAMA FARMERS COOPERATIVE, INC.

Decatur, Alabama 35601

FEEDING DIRECTIONS:

Sheep and Goat Feed is a 15% protein, high energy feed . It is formulated to provide the nutrients needed for growth and reproduction in both sheep and goats. This feed contains Decoquinatate which sheep

producers prefer as a preventative to coccidiosis. This feed contains supplemental minerals and vitamins, with no added copper and is safe for sheep. Feed to supplement good quality hay or pasture at a rate of 1 to 4 pounds per head per day depending on animal weight and stage of production. 1 to 4 pounds will provide not less than 7.5mg nor more than 30mg Decoquinatone per head per day. Always provide free choice minerals and vitamins for optimum intake. Always feed this product with hay or other forage. This feed is ideal for the maintenance and upkeep of mature goats and sheep. Nutrient requirements for nursing animals will increase and the feeding program should be changed to take this into account. Do not feed this product free choice to sheep or goats. Do not feed to horses or cattle.

Important Management Practices

Provide a source of clean fresh water and free choice minerals at all times. When making any change in an animals diet, plan for a 7 to 10 day transition period. Provide ample feeder space and do not allow feed to become wet or moldy. Poorly managed feeding programs increases the possibility of nutritional disorders. Consult an Animal Nutritionist if you have any questions or concerns.